

Superconducting Instability in a One-Dimensional Lattice of Berry Phase Molecules

D. Shelton and A. M. Tsvelik

February 1, 2008

*Department of Physics, University of Oxford, 1 Keble Road,
Oxford, OX1 3NP, UK*

Abstract

We study the one-dimensional model of conduction electrons interacting with local vibronic modes. States in the conduction band are two-fold degenerate both in orbital index and spin. It is shown that such one-dimensional system has a strong tendency for superconductivity. The superconducting pairing originates from coupling of two charge density waves; the scaling dimension of the superconducting order parameter is found to be close to $1/4$.

PACS numbers: 74.65.+n, 75.10. Jm, 75.25.+z

The discovery of superconductivity with fairly high transition temperatures in the alkali-intercalated fullerenes A_nC_{60} ($A = K, Rb$ and Cs) has revived an interest to the Jahn-Teller phenomenon. It is widely believed that the origin of superconducting pairing in the C_{60} compounds is due to the interaction of electrons with the vibronic modes of C_{60} molecules. A nontrivial aspect of this interaction is an orbital degeneracy of the local electronic states. In the strong coupling limit when one neglects the intermolecular electron hopping, each C_{60} molecule is described as a system of three local vibronic modes interacting with a three-fold degenerate electronic orbital. This interaction is non-diagonal thus leading to violations of the adiabatic approximation (the problem of Berry phase)

and, as a consequence, to a strong renormalization of the electronic wave functions (the Dynamical Jahn-Teller effect; see Ref. 1 for a review).

It is interesting to know whether a local degeneracy can survive in the presence of an intermolecular hopping and if yes, how it will affect the low-energy dynamics of the electronic subsystem. One obvious choice would be a structure where C_{60} molecules are situated on chains. In this case the one-dimensional electronic band is split into a single degenerate band created by the s, p_z orbitals and a doubly degenerate band created by p_x, p_y orbitals. The measurements performed for RbC_{60} give many indications of a quasi-one-dimensional behaviour². The direct band structure calculations done for A_nC_{60} -compounds with $A = K, Rb$ show, however, that the three-fold degeneracy is lifted with the characteristic energy of the order of $0.5eV$ and the resulting electronic bands have an essentially three dimensional character³. However, even if the results of the band theory calculations are correct, the C_{60} -based materials are not very far from being one-dimensional. The authors of Ref. 4 remark that the interchain hopping is very sensitive to the distance between the chains of C_{60} and therefore by alloying the alkali with “spacer molecules” such as NH_3 one can achieve a higher degree of one-dimensionality.

It is well known that in the standard one-dimensional attractive Hubbard model the superconductivity competes with the charge density wave instability. At small coupling constant the scaling dimension of the superconducting order parameter is close to 1. This is clearly seen for the the simplest exactly solvable model of fermions with a point-like interaction (see, for example, Ref. 4, where the relevant correlation functions for this model are listed). As we shall demonstrate in this paper, the superconducting pairing occurs in our model between electrons from different orbitals. Thus the presence of orbital symmetry is crucial in this respect. We shall also demonstrate that the scaling dimension of the order parameter is $1/4$, that is fairly small. A similar result was obtained for a one-dimensional lattice of Berry phase molecules in the strong coupling limit by Manini *et al.*⁵ and Santoro *et al.*⁶.

In this paper we study the model introduced in Ref. 5; it describes an one-dimensional chain of Jahn-Teller molecules with electrons belonging to two degenerate local orbitals hopping along the chain and interacting with local optical (vibronic) modes of the molecules. The Hamiltonian is given by

$$\begin{aligned}
H &= \sum_r \sum_{\alpha=\pm 1} \sum_{a=1,2} \left\{ -\frac{1}{2} t [c_{\alpha,a}^+(r+1) c_{\alpha,a}(r) + c_{\alpha,a}^+(r) c_{\alpha,a}(r+1)] \right. \\
&\quad \left. + J \left(c_{r,\alpha,a}^+ \hat{\sigma}_{ab}^x c_{r,\alpha,b} X_r + c_{r,\alpha,a}^+ \hat{\sigma}_{ab}^y c_{r,\alpha,b} Y_r \right) \right\} \\
&= \sum_r \left(-\frac{1}{2M} \frac{\partial^2}{\partial X_r^2} - \frac{1}{2M} \frac{\partial^2}{\partial Y_r^2} + \frac{k X_r^2}{2} + \frac{k Y_r^2}{2} \right) \quad (1)
\end{aligned}$$

Here $c_{\alpha,a}^+(r)$, $c_{\alpha,a}(r)$ are creation and annihilation of electrons on the site r and $n_a = \sum_{\alpha} c_{\alpha,a}^+ c_{\alpha,a}$. The Greek indices correspond to a spin projection and the English ones denote two degenerate orbitals. The electrons interact with local vibronic modes X_r , Y_r . As we have mentioned above, a single C_{60} molecule has a higher degeneracy, but we assume that such high degeneracy does not survive in a crystal.

Instead of the adiabatic approximation which is a standard tool in the theory of the Jahn-Teller effect we shall use another approach. Namely, we neglect all retardation effects related to the kinetic energy of vibrons and integrate over the vibronic modes. As in the standard BCS theory, this will give us a high energy cut-off of the order of the vibronic frequency $\omega_0 = (k/M)^{1/2}$. As the result we obtain the following effective interaction between the fermions:

$$V_{\text{int}} = -\frac{1}{2} g \sum_r [(c_r^+ \sigma^x c_r)^2 + (c_r^+ \sigma^y c_r)^2] = -g \sum_r c_{1,\alpha}^+ c_{1,\beta} c_{2,\beta}^+ c_{1,\alpha} \quad (2)$$

where $g = J^2/k$.

Assuming that $g/t \ll 1$ we linearize the electronic spectrum and then apply the standard bosonization procedure. At the first step we represent the fermionic operators as sums of right and left moving components

$$c_r = R(x) e^{-ik_F x} + L(x) e^{ik_F x} \quad (3)$$

where $x = ra$ (a is the lattice constant). Substituting this into Eq.(2) and keeping only non-oscillatory terms we get

$$V_{\text{int}} = V_1 + V_2 + V_3 \quad (4)$$

$$V_1 = g [R_{1,\alpha}^+ R_{1,\beta} L_{2,\beta}^+ L_{2,\alpha} + R_{2,\alpha}^+ R_{2,\beta} L_{1,\beta}^+ L_{1,\alpha}] \quad (5)$$

$$V_2 = -g [L_{1,\alpha}^+ R_{2,\beta}^+ R_{1,\beta} L_{2,\alpha} + L_{2,\alpha}^+ R_{1,\beta}^+ R_{2,\beta} L_{1,\alpha}] \quad (6)$$

$$V_3 = g [R_{1,\alpha}^+ R_{1,\beta} R_{2,\beta}^+ R_{2,\alpha} + (R \rightarrow L)] \quad (7)$$

We proceed further by bosonizing the fermionic operators:

$$R_{a,\sigma}, R_{a,\sigma}^+ = \frac{1}{\sqrt{2\pi a_0}} \exp(\pm i\varphi_{a,\sigma}),$$

$$L_{a,\sigma}, L_{a,\sigma}^+ = \frac{1}{\sqrt{2\pi a_0}} \exp(\mp i \bar{\varphi}_{-a,\sigma}) \quad (8)$$

$$R_{a,\sigma}^+ R_{a,\sigma} = \frac{1}{\pi} \partial_z \Phi_\sigma^a, \quad L_{a,\sigma}^+ L_{a,\sigma} = \frac{1}{\pi} \partial_{\bar{z}} \Phi_\sigma^a$$

$$\Phi(z, \bar{z}) = \varphi(z) + \bar{\varphi}(\bar{z}), \quad \Theta(z, \bar{z}) = \varphi(z) - \bar{\varphi}(\bar{z}) \quad (9)$$

where $a_0 = \Lambda^{-1}$ is the cut-off and $a = 1, 2$ is the orbital index, $\sigma = \pm 1$ is the spin index and $z = \tau + ix$, $\bar{z} = \tau - ix$. In order to simplify the Hamiltonian we introduce bosonic variables for charge (c) and spin (s) degrees of freedom on each orbital by

$$\varphi_{a,\sigma} = \frac{1}{\sqrt{2}} (\varphi_{a,c} + \sigma \varphi_{a,s}) \quad (10)$$

and then the combinations corresponding to total (+) and relative (-) charge and spin density fluctuations:

$$\varphi_{\pm,c,s} = \frac{1}{\sqrt{2}} (\varphi_{c,s}^1 \pm \varphi_{c,s}^2) \quad (11)$$

The bosonized Hamiltonian has the following form:

$$H = H_0 + V_1 + V_2 + V_3$$

$$H_0 = \frac{1}{2} \sum_{\pm} \left[v_c K_c^{\pm} (\partial_x \Theta_c^{\pm})^2 + \frac{v_c}{K_c^{\pm}} (\partial_x \Phi_c^{\pm})^2 + v_s K_s^{\pm} (\partial_x \Theta_s^{\pm})^2 + \frac{v_s}{K_s^{\pm}} (\partial_x \Phi_s^{\pm})^2 \right] \quad (12)$$

$$V_1 = \frac{g}{(\pi a_0)^2} \cos \Phi_s^+ \cos \Theta_s^-$$

$$V_2 = \frac{g}{(\pi a_0)^2} [\cos \Phi_c^- \cos \Phi_s^- + \cos \Phi_c^- \cos \Theta_s^-]$$

$$V_3 = \frac{g}{(\pi a_0)^2} \cos \Phi_s^- \cos \Theta_s^- \quad (13)$$

The bosonization of V_1 and V_3 in (7) also generates the following terms:

$$V_4 = \frac{g}{2\pi^2} [(\partial_x \Phi_c^+)^2 - (\partial_x \Phi_c^-)^2 + (\partial_x \Phi_s^+)^2 - (\partial_x \Phi_s^-)^2] \quad (14)$$

which renormalize the velocities $v_{c,s}$ and the exponents $K_{c,s}^{\pm}$ from their bare values to

$$\tilde{K}_{c,s}^{\pm} = (1 \pm g/\pi^2 v_{c,s})^{-1/2} K_{c,s}^{\pm}$$

They can therefore be absorbed into the bare Hamiltonian (12). The resulting Hamiltonian is a particular case of the Hamiltonian considered by Khveshchenko and Rice in relation to the problem of two coupled Hubbard chains⁷. Therefore we can use the renormalization group equations derived in this work. We simplify

these equations assuming that deviations of K 's from 1 are small ($K_{c,s}^{\pm} = 1 \mp x_c^{\pm}$) and expanding to the lowest nontrivial order in x :

$$\begin{aligned}\dot{g}_1 &= (2 - K_s^+ - \frac{1}{K_s^-})g_1 \approx (x_s^+ + x_s^-)g_1 \\ \dot{g}_2 &= (2 - K_c^- - K_s^-)g_2 \approx (x_c^- + x_s^-)g_2 \\ \dot{g}_3 &= (2 - K_c^- - \frac{1}{K_s^-})g_3 \approx (x_c^- - x_s^-)g_3\end{aligned}\tag{15}$$

where g_1 refers to the term V_1 and g_2 and g_3 to the first and the second term of V_2 respectively. The interaction V_3 is irrelevant. The dots stand for differentiation with respect to $\xi = \ln(\Lambda/\max(\omega, q))$. The equations for the exponents are

$$\begin{aligned}\dot{K}_c^+ &= 0, \quad \dot{K}_c^- = -\frac{1}{2}(K_c^-)^2(g_2^2 + g_3^2) \\ \dot{K}_s^+ &= -\frac{1}{2}(K_s^+)^2g_1^2, \quad \dot{K}_s^- = -\frac{1}{2}[(K_s^-)^2g_2^2 - g_1^2 - g_3^2]\end{aligned}\tag{16}$$

We note that the first equation is exact in all orders since the Hamiltonian is quadratic in the Φ_c^+ -field. To the first order in $x_{c,s}^{\pm}$ we have, defining $x_1 = (x_s^+ + x_s^-)$, $x_2 = (x_c^- + x_s^-)$, $x_3 = (x_c^- - x_s^-)$:

$$\dot{g}_i = x_i g_i\tag{17}$$

$$\begin{aligned}2\dot{x}_1 &= 2g_1^2 - g_2^2 + g_3^2, \quad 2\dot{x}_2 = 2g_2^2 - g_1^2 \\ 2\dot{x}_3 &= g_1^2 + 2g_3^2\end{aligned}\tag{18}$$

In order to determine the character of the renormalization group flow it is convenient to rewrite this system using the variables $y_i = \ln(g_i^2)$ such that $\dot{y}_i = 2x_i$. Then Eqs.(18) become

$$\begin{aligned}\ddot{y}_1 &= 2e^{2y_1} - e^{2y_2} + e^{2y_3} \\ \ddot{y}_2 &= 2e^{2y_2} - e^{2y_1} \\ \ddot{y}_3 &= e^{2y_1} + 2e^{2y_3}\end{aligned}\tag{19}$$

The right hand sides of all these equations never vanish simultaneously. Therefore we conclude that coupling constants grow. Thus we can be sure that the system scales to strong coupling.

From the behaviour of x_s^+ , $x_{c,s}^-$ we conclude that in the strong coupling regime gaps open in the spectrum of Φ_s^+ , $\Phi_{c,s}^-$ and the corresponding complex exponents acquire non-zero expectation values. From the form of the interaction (13) it is evident that non-zero expectation values are acquired by exponents of the fields

$\Phi_c^-, \Phi_s^+, \Theta_s^-$. The field Φ_c^+ remains a free field. This situation is very advantageous for the superconductivity. Indeed, the superconducting order parameter

$$\begin{aligned} \Delta = R_{1,+} L_{2,-} &\sim \exp \left[\frac{i}{2} (\Phi_s^- + \Theta_s^- + \Phi_s^+ + \Theta_c^+) \right] \\ &\sim m^2 : \exp \left(-\frac{i}{2} \Theta_c^+ \right) : \end{aligned} \quad (20)$$

and the asymptotics of the two-point correlation function at distances greater than m^{-1} is given by

$$\begin{aligned} \langle \Delta_a(1) \Delta_b^+(2) \rangle &\sim m^{3/2} [\tau_{12}^2 + x_{12}^2]^{-1/4K_c^+} \\ \frac{1}{K_c^+} &= \left(1 + \frac{g}{\pi^2 v_c} \right)^{1/2} \end{aligned} \quad (21)$$

At small g/v_c this function behaves almost like the correlation function of hard-core bosons (the transverse Pauli matrices in the XY-model). Since the latter model has the same central charge $c = 1$, we conclude that the gapless excitations in the present model are well described by the model of hard-core bosons. Thus our results are in complete agreement with the results of Santoro et al.⁶ obtained within a different approach. Such a small scaling dimension increases the probability of having a large transition temperature in a quasi-three-dimensional system built of weakly interacting chains. If t is the interchain Josephson coupling, then the dimensional analysis gives for the temperature of superconducting transition the estimate

$$T_c \sim m(t/m)^{\frac{1}{2-2d}}$$

In the conventional case $d \approx 1$ and the transition temperature is exponentially small in m/t , but for the present model $d = 1/4K_c^+$ and at small g/v_c we have $T_c \sim t^{2/3}$.

The authors are grateful to M. C. M. O'Brien and E. Tosatti for the interest to the work and G. Santoro for sending his preprint.

References

- [1] M. C. M. O'Brien and C. C. Chancey, Am. J. Phys. **61**, 688 (1993).
- [2] O. Chauvet, G. Oszlanyi, L. Forro, P. W. Stephens, M. Tegze, G. Faigel and A. Janossy, Phys. Rev. Lett. **72**, 2721 (1994).

- [3] S. C. Erwin, G. V. Krishna and E. J. Mele, sissa preprint (1994).
- [4] H. Frahm and V. Korepin, Phys. Rev. B **42**, 10553 (1990).
- [5] N. Manini, E. Tosatti and S. Doniach, to appear in Phys. Rev. B (1994).
- [6] G. Santoro, M. Airoidi, E. Tosatti and A. Parola, sissa preprint (1994).
- [7] V. Khveshchenko and M. Rice, Phys. Rev. B**50**, 252 (1994).